

Tests of scalar-tensor gravity*

Gilles Esposito-Farèse

*GRεCO, FRE 2435-CNRS, Institut d'Astrophysique de Paris,
98bis boulevard Arago, F-75014 Paris, France*

(Dated: September 20, 2004)

Abstract

The best motivated alternatives to general relativity are scalar-tensor theories, in which the gravitational interaction is mediated by one or several scalar fields together with the usual graviton. The analysis of their various experimental constraints allows us to understand better which features of the models have actually been tested, and to suggest new observations able to discriminate between them. This talk reviews three classes of constraints on such theories, which are qualitatively different from each other: (i) solar-system experiments; (ii) binary-pulsar tests and future detections of gravitational waves from inspiralling binaries; (iii) cosmological observations. While classes (i) and (ii) impose precise bounds respectively on the first and second derivatives of the matter-scalar coupling function, (iii) *a priori* allows us to reconstruct the full shapes of the functions of the scalar field defining the theory, but obviously with more uncertainties and/or more theoretical hypotheses needed. Simple arguments such as the absence of ghosts (to guarantee the stability of the field theory) nevertheless suffice to rule out a wide class of scalar-tensor models. Some of them can be probed only if one takes simultaneously into account solar-system and cosmological observations.

* Invited talk at the workshop *Phi in the Sky: The Quest for Cosmological Scalar Fields*, Porto, July 2004

I. INTRODUCTION

In the most natural alternative theories to general relativity (GR), gravity is mediated not only by a (spin-2) graviton corresponding to a metric $g_{\mu\nu}$, but also by a (spin-0) scalar field φ . Such scalar partners generically arise in all extra-dimensional theories, and notably in string theory. A dilaton is indeed already present in the supermultiplet of the 10-dimensional graviton, and several other scalar fields (called the moduli) also appear when performing a Kaluza-Klein dimensional reduction to our usual spacetime. They correspond to the components of the metric tensor g_{mn} in which m and n label extra dimensions. Moreover, contrary to other alternative theories of gravity, scalar-tensor theories respect most of GR's symmetries: conservation laws, constancy of non-gravitational constants, and local Lorentz invariance even if a subsystem is influenced by external masses. They can also satisfy exactly the weak equivalence principle (universality of free fall of laboratory-size objects) even for a massless scalar field.

Scalar fields are also involved in the cosmological models which reproduce most consistently present observational data. In particular, inflation theory is based on the presence of a scalar φ in a potential $V(\varphi)$ (for instance parabolic). It behaves as a fluid with a positive energy density $8\pi G\rho_\varphi = \dot{\varphi}^2 + 2V(\varphi)$ but a negative pressure $8\pi Gp_\varphi = \dot{\varphi}^2 - 2V(\varphi)$. This causes a period of exponential expansion of the universe, which can explain why causally disconnected regions at present may have been connected long ago. The isotropy of the observed cosmic microwave background (CMB) can thus be understood. Inflation also predicts that our universe is almost spatially flat, because any initial curvature has been exponentially reduced by the expansion. This is in remarkable agreement with the location of the first acoustic peak of the CMB spectrum at a multipolar index $\ell \simeq 220$ [1]. Various observations, notably of type Ia supernovae [2], tell us that there is about 70% of negative-pressure dark energy in our present universe ($\Omega_\Lambda \simeq 0.7$), suggesting that its expansion has been re-accelerating recently, since redshifts $z \sim 1$. This can be explained by the presence of a cosmological constant Λ in GR, but the quantity $\Omega_\Lambda \simeq 0.7$ translated in natural units gives an extremely small value $\Lambda \simeq 3 \times 10^{-122} c^3/(\hbar G)$, very problematic for particle physics if Λ is to be interpreted as the vacuum energy. This is the main reason why “quintessence” models have been proposed, in which the cosmological constant is replaced again by the potential $V(\varphi)$ of a scalar field. Its evolution towards a minimum of V during the cosmological expansion then explains more naturally why the present value $V(\varphi_0) \simeq \Lambda/2$ is so small.

Besides these theoretical and experimental reasons for studying scalar-tensor theories of gravity, one of their greatest interests is to embed GR within a class of mathematically consistent alternatives, in order to understand better which theoretical features have been experimentally tested, and which can be tested further. To simplify this review, we will restrict it to models involving a single scalar field, although the study of tensor-multi-scalar theories can also be done in great detail [3]. It suffices to note that their phenomenology is richer but similar to the single scalar case, at least when all scalar degrees of freedom carry positive energy, as required for the vacuum to be stable. In view of the very precise experimental verifications of the weak equivalence principle, we will also focus this review on theories which satisfy it exactly. In other words, we will assume that the action of geometry on matter is the same as in GR, but that the dynamics of geometry and the action of matter on it is modified because of the presence of a scalar field (see J.P. Uzan's contribution to the present proceedings, and in particular the upper-right panel of its Fig. 1). All matter fields, including gauge bosons, will thus be assumed to be universally coupled to one second rank

symmetric tensor, say $\tilde{g}_{\mu\nu}$. The difference with GR lies in the kinetic term of this tensor, which is now a mixing of spin-2 and spin-0 excitations. One indeed defines $\tilde{g}_{\mu\nu} \equiv A^2(\varphi)g_{\mu\nu}$, where $A(\varphi)$ is a function of the scalar field, while $g_{\mu\nu}$ denotes the Einstein (spin-2) metric. We will mainly consider the following action [4],

$$S = \frac{c^3}{4\pi G} \int \sqrt{-g} \left\{ \frac{R}{4} - \frac{1}{2}(\partial_\mu \varphi)^2 - V(\varphi) \right\} + S_{\text{matter}} [\text{matter}; \tilde{g}_{\mu\nu} \equiv A^2(\varphi)g_{\mu\nu}] , \quad (1)$$

which depends on two functions of the scalar field: the coupling function $A(\varphi)$ to matter, and a potential $V(\varphi)$. [Our signature is $-+++$, R is the scalar curvature of $g_{\mu\nu}$, and g its determinant.]

In Section II, we will review the main experimental constraints on this coupling function $A(\varphi)$, coming from two qualitatively different classes of data: solar-system experiments and binary-pulsar observations [5, 6]. In Section III, we will underline that cosmological observations give again very different types of constraints: Although they are more noisy, they *a priori* allow the reconstruction of both $A(\varphi)$ and $V(\varphi)$ [7, 8]. The most general scalar-tensor theories will be defined in Sec. IV, and we will illustrate in Sec. V, on a very special but interesting case, that one must sometimes take simultaneously into account solar-system and cosmological observations to obtain significant constraints on a theory [9]. We will finally give our conclusions in Sec. VI.

II. SOLAR-SYSTEM AND BINARY-PULSAR CONSTRAINTS

A. Solar-system tests

A massive scalar field has a negligible effect on the motion of celestial bodies if its mass is large with respect to the inverse of the interbody distances. On the other hand, if its mass is small enough, its potential $V(\varphi)$ can be locally neglected, but its coupling function to matter, $A(\varphi)$, is strongly constrained by experiment, as we will see below. The intermediate case, for which the size of the experiment is of the same order as the inverse scalar mass, is the main topic of E. Adelberger’s contribution to the present proceedings (see also [10]). The “chameleon” field [11] is another case where both the potential and the matter–scalar coupling function must be taken into account in solar-system experiments; see P. Brax’s contribution to these proceedings. In the present section, we will review the constraints on $A(\varphi)$ when the potential $V(\varphi)$ can be neglected.

The precision of solar-system observations allow us not only to test Newton’s law, but also its relativistic corrections $\propto 1/c^2$, called “post-Newtonian”. At this order, the predictions of metric theories of gravity can be parametrized by a set of 10 real numbers in the so called “PPN” formalism (parametrized post-Newtonian) [12]. All of them are presently constrained to be very close to their general relativistic values [13], and in particular the two famous parameters β^{PPN} and γ^{PPN} introduced by Eddington in the Schwarzschild metric:

$$\begin{aligned} -g_{00} &= 1 - 2\frac{Gm}{rc^2} + 2\beta^{\text{PPN}} \left(\frac{Gm}{rc^2} \right)^2 + \mathcal{O} \left(\frac{1}{c^6} \right) , \\ g_{ij} &= \delta_{ij} \left(1 + 2\gamma^{\text{PPN}} \frac{Gm}{rc^2} \right) + \mathcal{O} \left(\frac{1}{c^4} \right) . \end{aligned} \quad (2)$$




FIG. 1: Solar-system and binary-pulsar constraints on the matter-scalar coupling function $\ln A(\varphi) = \alpha_0(\varphi - \varphi_0) + \frac{1}{2}\beta_0(\varphi - \varphi_0)^2 + \mathcal{O}(\varphi - \varphi_0)^3$. The allowed region is shaded. The vertical axis ($\beta_0 = 0$) corresponds to Brans-Dicke theory with a parameter $2\omega_{\text{BD}} + 3 = 1/\alpha_0^2$. The horizontal axis ($\alpha_0 = 0$) corresponds to theories which are perturbatively equivalent to GR, i.e., which predict strictly no deviation from it (at any order $1/c^n$) in the weak-field conditions of the solar system.

General relativity corresponds to $\beta^{\text{PPN}} = \gamma^{\text{PPN}} = 1$. In scalar-tensor theories [3, 5], these two latter parameters are the only ones which can differ from their GR values. They are related to the first two derivatives of

$$\ln A(\varphi) \equiv \alpha_0(\varphi - \varphi_0) + \frac{1}{2}\beta_0(\varphi - \varphi_0)^2 + \mathcal{O}(\varphi - \varphi_0)^3, \quad (3)$$

computed at the background value φ_0 of the scalar field:

$$\gamma^{\text{PPN}} - 1 = -\frac{2\alpha_0^2}{1 + \alpha_0^2}, \quad \beta^{\text{PPN}} - 1 = \frac{1}{2} \frac{\alpha_0\beta_0\alpha_0}{(1 + \alpha_0^2)^2}. \quad (4)$$

The factor α_0^2 comes from the exchange of a scalar particle between two bodies, whereas $\alpha_0\beta_0\alpha_0$ comes from a scalar exchange between three bodies. Solar system experiments impose the constraints displayed as thin lines in Fig. 1. The Mercury symbol refers to the perihelion shift of this planet, whose observed value imply the bound [14]

$$|2\gamma^{\text{PPN}} - \beta^{\text{PPN}} - 1| < 3 \times 10^{-3}, \quad (5)$$

the Moon symbol refers to Lunar Laser Ranging [15]:

$$4\beta^{\text{PPN}} - \gamma^{\text{PPN}} - 3 = (-0.7 \pm 1) \times 10^{-3}, \quad (6)$$




FIG. 2: Heuristic argument to explain the phenomenon of “spontaneous scalarization”. When $\beta_0 < 0$ and the compactness Gm/Rc^2 of a body is large enough, it is energetically favorable to create a local scalar field different from the background value. The body becomes thus strongly coupled to the scalar field.

the star symbol to light deflection as measured by Very Long Baseline Interferometry [16]:

$$|\gamma^{\text{PPN}} - 1| < 4 \times 10^{-4}, \quad (7)$$

and the label “Cassini” to the impressive recent constraint obtained by measuring the time delay variation to the Cassini spacecraft near solar conjunction [17]:

$$\gamma^{\text{PPN}} - 1 = (2.1 \pm 2.3) \times 10^{-5}. \quad (8)$$

Solar-system tests thus constrain the first derivative α_0 to be small, i.e., the linear interaction between matter and the scalar field to be weak. On the other hand, the second derivative β_0 may take large values (of either sign), i.e., matter may be strongly coupled to two scalar lines.

B. Nonperturbative strong-field effects

At higher post-Newtonian orders $1/c^n$, a simple diagrammatic argument shows that any deviation from GR involves at least two factors α_0 , and has the schematic form

$$\text{deviation from GR} = \alpha_0^2 \times \left[\lambda_0 + \lambda_1 \frac{Gm}{Rc^2} + \lambda_2 \left(\frac{Gm}{Rc^2} \right)^2 + \dots \right], \quad (9)$$

where m and R denote the mass and radius of the considered body, and $\lambda_0, \lambda_1, \dots$ are constants built from the coefficients $\alpha_0, \beta_0, \dots$ of expansion (3). Since α_0^2 is experimentally known to be small, we thus expect the theory to be close to GR at any order. However, some nonperturbative effects may occur in strong-field conditions: If the compactness Gm/Rc^2




FIG. 3: Scalar charge α_A versus baryonic mass $\bar{m}_A$, for the model $A(\varphi) = \exp(-3\varphi^2)$ (i.e., $\beta_0 = -6$). The solid line corresponds to the maximum value of α_0 allowed by Eqs. (5)-(6), and the dashed line to $\alpha_0 = 0$. The dotted lines correspond to unstable configurations of the star.

of a body is greater than a critical value, the square brackets of Eq. (9) can become large enough to compensate even a vanishingly small α_0^2 . This can happen notably for neutron stars, whose compactnesses are of order $Gm/Rc^2 \sim 0.2$, as compared to only 2×10^{-6} for the Sun or even 7×10^{-10} for the Earth. To illustrate this, let us consider a model for which α_0 vanishes strictly, i.e., which is perturbatively equivalent to GR: There is strictly no deviation from GR at any order in a perturbative expansion in powers of $1/c$. A parabolic coupling function $\ln A(\varphi) = \frac{1}{2}\beta_0\varphi^2$ suffices for our purpose (we set here $\varphi_0 = 0$ to simplify). At the center of a static body, the scalar field takes a particular value φ_c , and it decreases as $1/r$ outside. The energy of such a scalar field configuration involves two contributions, coming respectively from the kinetic term and from the matter-scalar coupling function in action (1). As a rough estimate of its value, one can write

$$\text{Energy} \simeq \int \left[\frac{1}{2}(\partial_i \varphi)^2 + \rho e^{\beta_0 \varphi^2/2} \right] \simeq mc^2 \left(\frac{\varphi_c^2/2}{Gm/Rc^2} + e^{\beta_0 \varphi_c^2/2} \right). \quad (10)$$

When $\beta_0 < 0$, this is the sum of a parabola and a Gaussian, and if the compactness Gm/Rc^2 is large enough, the function $\text{Energy}(\varphi_c)$ has the shape of a Mexican hat, see Fig. 2. The value $\varphi_c = 0$ now corresponds to a local *maximum* of the energy. It is therefore energetically favorable for the star to create a nonvanishing scalar field φ_c , and thereby a nonvanishing “scalar charge” $d \ln A(\varphi_c)/d\varphi_c = \beta_0 \varphi_c$. This phenomenon is analogous to the spontaneous magnetization of ferromagnets.

This heuristic argument has been verified by explicit numerical calculations, taking into account the coupled differential equations of the metric and the scalar field, and using various realistic equations of state to describe nuclear matter inside a neutron star [5]. The correct definition of the linear coupling strength between a compact body A and the scalar field reads $\alpha_A \equiv \partial \ln m_A / \partial \varphi_0$. It is plotted in Fig. 3 for the particular model $\beta_0 = -6$. One finds that there exists indeed a “spontaneous scalarization” above a critical mass (whose value decreases as $-\beta_0$ grows).

On the other hand, if $\beta_0 > 0$, both the above heuristic argument and the actual numerical calculations show that $|\alpha_A| < |\alpha_0|$. In that case, one finds that neutron stars are even less coupled to the scalar field than solar-system bodies. This decoupling is similar to the behavior of the “chameleon” field [11], but it is here caused by the nonlinearity of $\ln A(\varphi)$,




FIG. 4: Mass plane (m_A = pulsar, m_B = companion) of the Hulse-Taylor binary pulsar PSR B1913+16 in general relativity (left panel) and for a scalar-tensor theory with $\beta_0 = -6$ (right panel). The widths of the lines are larger than 1σ error bars. While GR passes the test with flying colors, the value $\beta_0 = -6$ is ruled out.

whereas the chameleon's decoupling is due to a competition between $A(\varphi)$ and the potential $V(\varphi)$.

C. Binary-pulsar tests

The scalar charge α_A enters the predictions of the theory in the same way as α_0 in weak-field conditions. For instance, in the orbital motion of two bodies A and B, the Eddington parameter γ^{PPN} keeps the form of Eq. (4), but it now involves the product $\alpha_A \alpha_B$ of the two scalar charges instead of α_0^2 . Similarly, the strong-field analogue of β^{PPN} involves products of scalar charges and their derivatives $\beta_A \equiv \partial \alpha_A / \partial \varphi_0$. Since $|\alpha_A| \simeq 0.6$ in the model of Fig. 3, one thus expects deviations by $\sim 35\%$ from some general relativistic predictions. Moreover, the quadratic coupling strength β_A can take very large numerical values near the critical mass, like the magnetic susceptibility of ferromagnets. Therefore, even larger deviations from GR are found when the mass of a neutron star happens to be close to the critical one.

Binary pulsars are ideal tools for testing gravity in strong-field conditions. A pulsar is a rapidly rotating neutron star emitting a beam of radio waves, like a lighthouse. Experiment tells us that isolated pulsars are very stable clocks, when they are old enough. A pulsar A orbiting a companion B is thus a moving clock, the best tool that one could dream of to test a relativistic theory. Indeed, by precisely timing its pulse arrivals, one gets a stroboscopic information on its orbit, and one can measure several relativistic effects. Such effects do depend on the two masses m_A , m_B , which are not directly measurable. However, two different effects suffice to determine them, and a third relativistic observable then gives a test of the theory. In the case of the famous Hulse-Taylor binary pulsar PSR B1913+16, three relativistic parameters have been determined with great accuracy [18]: (i) the Einstein time delay parameter γ_T , which combines the second-order Doppler effect ($\propto v_A^2/2c^2$, where v_A is the pulsar's velocity) together with the redshift due to the companion ($\propto Gm_B/r_{AB}c^2$,

where r_{AB} the pulsar-companion distance); (ii) the periastron advance $\dot{\omega}$ (relativistic effect of order v^2/c^2); and (iii) the rate of change of the orbital period, $\dot{P}$, caused by gravitational radiation damping (an effect of order v^5/c^5 in GR, but of order v^3/c^3 in scalar-tensor theories; see below). Figure 4 displays the plane of the two *a priori* unknown masses m_A and m_B . For each relativistic parameter, the prediction of a given theory is consistent with the observed value only along a thin line. In GR, the fact that the three lines meet in one point means that there exists a pair of masses (m_A, m_B) simultaneously consistent with all three observables. This is thus a spectacular confirmation of Einstein's theory.

Obviously, these lines are deformed in scalar-tensor theories, and if they do not have any common intersection, the corresponding model is ruled out. The right panel of Fig. 4 illustrates such a case. The allowed theories lie below and to the right of the line labeled PSR B1913+16 in Fig. 1. This plot shows vividly the qualitative difference between solar-system and binary-pulsar observations. Indeed, the latter impose

$$\beta_0 > -4.5, \quad (11)$$

even for a vanishingly small α_0 . This constraint is due to the spontaneous scalarization of neutron stars, which occurs when $-\beta_0$ is large enough. Equations (4) allow us to rewrite this inequality in terms of the Eddington parameters β^{PPN} and γ^{PPN} , which are both consistent with 1 in the solar system. One finds

$$\frac{\beta^{\text{PPN}} - 1}{\gamma^{\text{PPN}} - 1} < 1.1. \quad (12)$$

The singular (0/0) nature of this ratio underlines why such a conclusion could not be obtained in weak-field experiments.

Several other binary pulsars are presently known and accurately timed. To simplify Fig. 1, we did not display all the constraints they impose in the theory plane (see Ref. [6]). We just plotted as a dotted line the tight ones imposed by the recently timed neutron star-white dwarf binary PSR J1141-6545 [19]. Contrary to neutron stars, the white dwarf companion is never spontaneously scalarized, because its compactness Gm/Rc^2 is too small. This binary system is thus very asymmetrical, and the pulsar's scalar charge α_A is generically much greater than its companion's, α_B . This causes a huge deviation from GR in the orbital period variation $\dot{P}$. Indeed, the energy flux carried out by gravitational waves is of the form

$$\begin{aligned} \text{Energy flux} = & \left\{ \frac{\text{Quadrupole}}{c^5} + \mathcal{O}\left(\frac{1}{c^7}\right) \right\}_{\text{helicity } 2} \\ & + \left\{ \frac{\text{Monopole}}{c} \left(0 + \frac{1}{c^2}\right)^2 + \frac{\text{Dipole}}{c^3} (\alpha_A - \alpha_B)^2 + \frac{\text{Quadrupole}}{c^5} + \mathcal{O}\left(\frac{1}{c^7}\right) \right\}_{\text{helicity } 0} \end{aligned} \quad (13)$$

The first curly brackets contain the prediction of general relativity, of order v^5/c^5 , whereas the second ones contain the extra contributions predicted in tensor-scalar theories. In particular, the dipolar contribution is of order v^3/c^3 , much larger than the usual quadrupole of GR if the two scalar charge are significantly different. This is the reason why the asymmetrical system PSR J1141-6545 is so constraining for scalar-tensor theories, although its experimental uncertainties on $\dot{P}$ are still rather large. Notice that this binary pulsar is almost as constraining as solar-system tests even in the region $\beta_0 > 0$; see Fig. 1. It should

probe values of the Eddington parameter $|\gamma^{\text{PPN}} - 1| \sim 10^{-6}$ by the end of the decade, i.e., an order of magnitude better than present solar-system limits (8).

The LIGO and VIRGO interferometers will detect gravitational waves emitted by inspiralling binaries. Since matter is known to be weakly coupled to the scalar field in the solar system (small value of $|\alpha_0|$), these detectors will not be sensitive of the helicity-0 waves. On the other hand, the time evolution of the helicity-2 chirp does depend on the energy flux (13), which differs significantly from the GR prediction when the scalar charges α_A , α_B are nonzero. Therefore, the GR wave templates used for matched filtering in LIGO and VIRGO may not be accurate if there exists a scalar partner to the graviton, and the signal-to-noise ratio may then drop. Fortunately, it was shown in [5] that binary-pulsar data are so precise that they already exclude the models which would have predicted significant effects in the gravitational waveforms. Therefore, although these interferometers are *a priori* more sensitive to the scalar field than classic solar-system tests, one may securely use the GR wave templates for their data analysis. On the other hand, it was shown in [20] that the LISA space interferometer can be sensitive to scalar effects which are still allowed by all present tests. Future binary-pulsar data should nevertheless probe them before the LISA mission is launched [6].

In conclusion, solar-system tests tightly constrain the first derivative of $\ln A(\varphi)$ (linear matter-scalar coupling strength α_0), whereas binary-pulsar data impose that its second derivative β_0 (quadratic coupling matter-scalar-scalar) is not large and negative. We will now see that cosmological observations give access to the full shape of this coupling function, of course not with the same accuracy as the above tests, but with the capability of constraining any higher derivative of $\ln A(\varphi)$ (vertex of matter with any number of scalar lines). Moreover, cosmological data can also give access to the full shape of the potential $V(\varphi)$.

III. RECONSTRUCTION OF A SCALAR-TENSOR THEORY FROM COSMOLOGICAL OBSERVATIONS

In cosmology, the usual approach to study quintessence models is to assume a particular form for the potential $V(\varphi)$ (and the matter-scalar coupling function $A(\varphi)$ when one considers “extended quintessence” models), to compute all possible observable predictions, and to compare them to experimental data.

In contrast, in the phenomenological approach, one wishes to *reconstruct* the Lagrangian of the theory from cosmological observations. It was proved in [7] that the knowledge of the luminosity distance $D_L(z)$ and of the density fluctuations $\delta_m(z) = \delta\rho/\rho$ as functions of the redshift z indeed suffices to reconstruct both the potential $V(\varphi)$ and the coupling function $A(\varphi)$. [The knowledge of the Hubble constant H_0 or the present matter density $\Omega_{m,0}$ is not even necessary, as they derive from $D_L(z)$ and $\delta_m(z)$.] In order to write the corresponding equations as simply as possible, it is convenient to perform a change of variables on action (1). Let us define

$$\begin{aligned} \tilde{g}_{\mu\nu} &\equiv A^2(\varphi)g_{\mu\nu}; & \Phi &\equiv A^{-2}(\varphi); \\ 2\omega_{\text{BD}}(\Phi) + 3 &\equiv A^2(\varphi)/A'^2(\varphi); & U(\Phi) &\equiv 2V(\varphi)/A^4(\varphi). \end{aligned} \quad (14)$$

Then action (1) takes the “Brans-Dicke” form

$$S = \frac{c^3}{16\pi G} \int d^4x \sqrt{-\tilde{g}} \left\{ \Phi \tilde{R} - \frac{\omega_{\text{BD}}(\Phi)}{\Phi} (\partial_\mu \Phi)^2 - 2U(\Phi) \right\} + S_{\text{matter}}[\text{matter}; \tilde{g}_{\mu\nu}], \quad (15)$$

in which the scalar curvature $\tilde{R}$, the determinant $\tilde{g}$, and the contraction of indices in $(\partial_\mu \Phi)^2$ now all correspond to the physical (“Jordan-frame”) metric $\tilde{g}_{\mu\nu}$. Note that this action defines exactly the *same* theory as Eq. (1).

The reconstruction equations can then be written straightforwardly. The first step is to derive the Hubble function $H(z)$ from the knowledge of the luminosity distance $D_L(z)$:

$$\frac{1}{H(z)} = \left(\frac{D_L(z)}{1+z} \right)' \times \left[1 + \Omega_{\kappa,0} \left(\frac{H_0 D_L(z)}{1+z} \right)^2 \right]^{-1/2}, \quad (16)$$

where a prime denotes here a derivation with respect to the redshift z , and $\Omega_{\kappa,0}$ is the possible (small) contribution of the Universe’s spatial curvature to the total energy density. In the short wavelength limit $\lambda \ll (H^{-1}, m_\varphi^{-1})$, the field equations for small perturbations yield $\ddot{\delta}_m + 2H\dot{\delta}_m - 4\pi G_{\text{eff}}\rho\delta_m \simeq 0$, where $G_{\text{eff}} \equiv G[A^2(\varphi) + (dA/d\varphi)^2]$ is the effective gravitational constant between two close test masses, taking into account the scalar-field contribution. In terms of the Brans-Dicke variables, this gives

$$\frac{\Phi}{\Phi_0} \simeq \frac{3}{2} \left(\frac{H_0}{H} \right)^2 \frac{(1+z) \Omega_{m,0} \delta_m}{\delta_m'' + \left(\frac{H'}{H} - \frac{1}{1+z} \right) \delta_m'} \times \left(1 + \frac{1}{2\omega_{\text{BD}} + 3} \right). \quad (17)$$

On the other hand, the background evolution equations yield

$$\begin{aligned} \frac{2U}{(1+z)^2 H^2} = & \Phi'' + \left(\frac{H'}{H} - \frac{4}{1+z} \right) \Phi' + \left[\frac{6}{(1+z)^2} - \frac{2}{1+z} \frac{H'}{H} - 4 \left(\frac{H_0}{H} \right)^2 \Omega_{\kappa,0} \right] \Phi \\ & - 3(1+z) \left(\frac{H_0}{H} \right)^2 \Phi_0 \Omega_{m,0}, \end{aligned} \quad (18)$$

$$\begin{aligned} \omega_{\text{BD}} = & -\frac{\Phi}{\Phi^2} \left\{ \Phi'' + \left(\frac{H'}{H} + \frac{2}{1+z} \right) \Phi' - 2 \left[\frac{1}{1+z} \frac{H'}{H} - \left(\frac{H_0}{H} \right)^2 \Omega_{\kappa,0} \right] \Phi \right. \\ & \left. + 3(1+z) \left(\frac{H_0}{H} \right)^2 \Phi_0 \Omega_{m,0} \right\}. \end{aligned} \quad (19)$$

Therefore, Eqs. (17)–(19) allow us to reconstruct $\Phi(z)$, $U(z)$ and $\omega_{\text{BD}}(z)$ as functions of the redshift, and thereby $U(\Phi)$ and $\omega_{\text{BD}}(\Phi)$ in a parametric way. Actually, the last factor of Eq. (17) may be neglected, since we know that $(2\omega_{\text{BD}} + 3)^{-1} = \alpha_0^2 \ll 1$ at present. In that case, Eqs. (17)–(19) become merely algebraic, and no integration of differential equation is required. However, one should note that second derivatives of experimental data are needed, so that large uncertainties are expected in such a reconstruction program (see R. Jimenez’s contribution to the present proceedings). The point that we wish to emphasize is that the full reconstruction of the microscopic action (15) is theoretically possible. Although it needs some algebra, this result seems anyway obvious: It is possible to *fit* two observed functions $[D_L(z)$ and $\delta_m(z)]$ thanks to two unknown ones $[V(\varphi)$ and $A(\varphi)$, or equivalently $U(\Phi)$ and $\omega_{\text{BD}}(\Phi)]$.

However, future experiments (like the SNAP satellite) will only give access to the luminosity distance $D_L(z)$ with a good accuracy, and the density contrast $\delta_m(z)$ cannot yet be used to constrain the models. A *semi-phenomenological* approach can thus be useful: We make some theoretical hypotheses on either the potential $V(\varphi)$ or the coupling function




FIG. 5: Matter-scalar coupling function $\ln A(\varphi)$ which reproduces exactly the same accelerated expansion as GR plus a cosmological constant (with $\Omega_\Lambda \simeq 0.7$), without any scalar field potential $V(\varphi)$ nor any cosmological constant. However, this model works only for redshifts $z \lesssim 0.7$.

$A(\varphi)$, and we reconstruct the other one from $D_L(z)$. *A priori*, one may think that such a reconstruction is again obvious: We fit one observed function $[D_L(z)]$ with one unknown function $[V(\varphi)$ or $A(\varphi)]$. But this naive reasoning is only valid locally, on a small interval. Indeed, the reconstructed function may for instance diverge for some value of the redshift, or one of the degrees of freedom may need to take a negative energy beyond a given redshift, which would make the theory unstable (and ill defined as a field theory on the surface where the energy changes its sign). The positivity of the graviton energy implies $A^2(\varphi) > 0$, which can be translated in terms of the Brans-Dicke scalar field as $\Phi > 0$. On the other hand, the positivity of the scalar-field energy imposes the minus sign in front of the scalar kinetic term $-(\partial_\mu \varphi)^2$ in action (1), which translates as $\omega_{\text{BD}} > -\frac{3}{2}$ in terms of the Brans-Dicke parameter. It was shown in Ref. [8] that these conditions impose tight constraints on the theories as soon as one knows $D_L(z)$ over a wide enough interval $z \in [0, \sim 2]$.

For instance, the present accelerated expansion of the universe can be perfectly described by a scalar-tensor theory with a vanishing potential $V(\varphi) = 0$ (and therefore a vanishing cosmological constant too). Reference [8] derived analytically the coupling function $A(\varphi)$ which reproduces exactly the same evolution of the scale factor $a(z)$ as the one predicted by GR plus a cosmological constant. Figure 5 displays this reconstructed function $\ln A(\varphi)$, which has a nice parabolic shape, with a minimum very close to the present value φ_0 of the scalar field, and a positive second derivative. This is not only consistent with binary-pulsar data [which forbid large and negative values of this second derivative β_0 , see Eq. (11)] but also with the cosmological attractor phenomenon analyzed by Damour and Nordtvedt [21]: The scalar field is generically attracted towards a minimum of $\ln A(\varphi)$ during the cosmological expansion, whereas some fine tuning would be necessary to reach a maximum (negative β_0). Therefore, we are in the difficult situation in which two very different theories are both consistent with experimental data, and there seems to be no way to distinguish them. Fortunately, Ref. [8] found that this scalar-tensor theory cannot mimic GR plus a cosmological constant beyond a redshift $z \sim 0.7$, because the scalar field φ would diverge at this value, and above all because the graviton energy would become negative beyond. Therefore, it suffices to measure $D_L(z)$ precisely enough up to $z \sim 1$ to rule out such a potential-free scalar-tensor theory. Actually, if $D_L(z)$ is measured over a wider interval $z \in [0, \sim 2]$, large

experimental errors (tens of percent) are not problematic: It is still possible to distinguish this potential-free model from GR plus a cosmological constant, and thereby to rule out one of them. The results of the SNAP satellite up to $z \sim 2$ will therefore be very useful to constrain scalar-tensor theories of gravity.

One can also impose a particular form of the coupling function $A(\varphi)$ and reconstruct the potential $V(\varphi)$ which reproduces the observed luminosity distance $D_L(z)$. For instance, for a minimally coupled scalar field $A(\varphi) = 1$ (usual “quintessence”) in a spatially curved universe, Ref. [8] analytically derived the expression of $V(\varphi)$ which gives the same cosmological evolution as GR plus a cosmological constant in a spatially flat universe. It was found that the shape of the potential is smoother when the universe is (marginally) closed. If it is flat or almost flat, one obviously recovers a cosmological constant with its unnaturally small value $\Lambda \simeq 3 \times 10^{-122} c^3 / (\hbar G)$. Therefore, in that case, aesthetic reasons may help us discriminate between the theories, instead of the much stronger argument of the positivity of energy that was used above. This shows anyway that the sole knowledge of $D_L(z)$ suffices to constrain scalar-tensor theories of gravity.

IV. THE MOST GENERAL SCALAR-TENSOR THEORIES

More general scalar-tensor theories than action (1) can be defined. First, one may consider several scalar fields [3], labeled by latin indices, φ^a . In the action, their kinetic term $-\frac{1}{2}\partial_\mu\varphi^a\partial^\mu\varphi^b$ will thus involve a matrix γ_{ab} to contract the indices. Such a matrix may itself depend on the scalar field, and it defines thus what is called a σ -model. An example of a tensor-bi-scalar theory is provided by hybrid inflation. Note that the σ -model metric γ_{ab} must be positive definite for all the scalar fields to carry positive energy.

The quintessence potential $V(\varphi^a)$ may also depend on the various scalar fields, and thereby define n -dimensional mountains along which the scalar degrees of freedom roll down. Similarly, the matter-scalar coupling function $A(\varphi^a)$ may have a nontrivial shape, and define a rich “extended quintessence” phenomenology.

If the weak equivalence principle is not exactly satisfied, different species of matter, labeled i , may have different coupling functions $A_i(\varphi^a)$ to the scalar field. It should be noted that superstring theory does predict such couplings, which violate equivalence principle tests by 9 or 10 orders of magnitude at tree level (see J.P. Uzan’s contribution to the present proceedings). To reconcile it with experiment, the standard argument is that no symmetry protects the masses of the scalar fields, so that all of them should acquire a large enough mass at present to be exponentially suppressed, or anyway a potential such that they are very weakly coupled to matter on Earth (cf. the chameleon model [11]). However, “flat directions” generically arise in low-energy effective models inspired by superstrings, i.e., one or several scalar fields which remain strictly massless and violate *a priori* the weak equivalence principle. In that case, string loop corrections can induce an attractor mechanism [25], which efficiently drives such scalars towards an extremum of their coupling functions to matter, so that they almost decouple at present.

It is also possible to couple the scalar fields to the Gauss-Bonnet topological invariant, $R^2_{\mu\nu\rho\sigma} - 4R^2_{\mu\nu} + R^2$. This is the only combination involving powers of the Riemann and/or the Ricci tensors which preserves the spectrum of the theory, i.e., which does not add any extra degree of freedom. A function $f(R)$ of the scalar curvature is well known to be equivalent to an extra scalar degree of freedom in the model, cf. [22]. It suffices to rewrite $\int f(R)$ as $\int [f(\varphi) + (R - \varphi)f'(\varphi)]$ and to show that the field equations deriving from these actions

coincide. The latter is of the Brans-Dicke type (15), with $\omega_{\text{BD}} = 0$, and a mere change of variables (conformal transformation) $g_{\mu\nu}^* \equiv f'(\varphi)g_{\mu\nu}$ yields the canonical form (1). Similarly, a function $f(R, \square R, \dots, \square^n R)$ of the scalar curvature and its iterated d'Alembertian can be shown to be equivalent to the introduction of $n + 1$ scalar fields in the theory [23]. On the other hand, functions $f(R_{\mu\nu})$ or $f(R_{\mu\nu\rho\sigma})$ of the Ricci or the Riemann tensor have been proved several times [24] to involve generically an extra massive spin-2 ghost, i.e., an extra negative-energy graviton which spoils the stability of the theory. The only allowed combination is the Gauss-Bonnet topological invariant.

Finally, one may also consider a function $F(\partial_\mu\varphi\partial^\mu\varphi)$ of the kinetic terms, as in “k-essence” models [26]. However, only some particular functions F preserve the positivity of the scalars’ kinetic energy.

In conclusion, the most general tensor–multi-scalar theories that we may consider are defined by the following action:

$$\begin{aligned}
S = & \frac{c^3}{4\pi G} \int \sqrt{-g} \left\{ \frac{R}{4} - \frac{1}{2} F\left(g^{\mu\nu}\gamma_{ab}(\varphi^c) \partial_\mu\varphi^a \partial_\nu\varphi^b\right) - V(\varphi^a) \right\} \\
& - \hbar \int \sqrt{-g} W(\varphi^a) (R_{\mu\nu\rho\sigma}^2 - 4R_{\mu\nu}^2 + R^2) \\
& + \sum_i S_{\text{matter}_i} [\text{matter}_i; \tilde{g}_{\mu\nu}^{(i)} \equiv A_i^2(\varphi^a)g_{\mu\nu}].
\end{aligned} \tag{20}$$

V. EXPERIMENTAL CONSTRAINTS ON A SCALAR–GAUSS-BONNET COUPLING

In this section, we will consider the very particular case of a scalar-tensor theory with a single scalar field φ , no matter-scalar coupling [$A(\varphi) = 1$], no potential [$V(\varphi) = 0$], and a standard kinetic term $-\frac{1}{2}(\partial_\mu\varphi)^2$, but with a scalar–Gauss-Bonnet coupling $W(\varphi)$:

$$\begin{aligned}
S = & \frac{c^3}{4\pi G} \int \sqrt{-g} \left\{ \frac{R}{4} - \frac{1}{2} (\partial_\mu\varphi)^2 \right\} - \hbar \int \sqrt{-g} W(\varphi) (R_{\mu\nu\rho\sigma}^2 - 4R_{\mu\nu}^2 + R^2) \\
& + S_{\text{matter}} [\text{matter}; g_{\mu\nu}].
\end{aligned} \tag{21}$$

Our aim is just to illustrate the kind of constraints which can be imposed on such a scalar–Gauss-Bonnet coupling [9]. This model will underline that solar-system and cosmological observations must be sometimes simultaneously taken into account to constrain a theory. The perturbative but highly nonlinear effects which happen can also be instructive.

Solar-system and binary-pulsar tests are local, and any deviation from GR depends on the magnitude of the scalar field created by a massive body. Let us thus analyze first the equation satisfied by φ in the vicinity of a spherical mass $m_\odot$. We can assume that the metric is close to the Schwarzschild solution, and we get at the first nonvanishing order in powers of $Gm_\odot/c^2$

$$\square\varphi = \frac{3\ell_0^2}{r^6} \left(\frac{2Gm_\odot}{c^2} \right)^2 [W'_0 + W''_0\varphi + \mathcal{O}(\varphi^2)] , \tag{22}$$

where we have set $\ell_0^2 \equiv 16\pi G\hbar/c^3$, and where the derivative $W'(\varphi)$ has been expanded in powers of φ in the right-hand side. Since we are assuming that φ takes small values, let




FIG. 6: Scalar-Gauss-Bonnet coupling function $W(\varphi)$ which exactly reproduces the cosmological expansion predicted by GR plus a cosmological constant.

us neglect the contribution $W_0''\varphi$. We can then compute any observable prediction, but we quote below only the results for the light deflection angle ($\Delta\theta_*$) and for the perihelion shift per orbit ($\Delta\theta_p$), which suffice for our purpose:

$$\Delta\theta_* = \frac{4Gm_\odot}{r_0c^2} + \frac{1536}{35} \left(\frac{Gm_\odot}{r_0c^2} \right)^3 \left(\frac{\ell_0}{r_0} \right)^4 W_0'^2, \quad (23)$$

$$\Delta\theta_p = \frac{6\pi Gm_\odot}{pc^2} + 192\pi \left(\frac{Gm_\odot}{pc^2} \right)^2 \left(\frac{\ell_0}{p} \right)^4 W_0'^2, \quad (24)$$

where r_0 is the minimal distance between the light ray and the Sun, and p is the *semilatus rectum* of an orbit. The first terms on the right-hand sides are the usual GR predictions, at first order in $Gm_\odot/c^2$. In conclusion, solar-system (and binary-pulsar) tests can easily be passed if $|W_0'|$ is small enough.

One can now reconstruct the full shape of $W(\varphi)$ from the cosmological observation of the luminosity distance $D_L(z)$, as in Sec. III above. We find that this can always been done, without any problem of negative energy, contrary to what happened above for the matter-scalar coupling function $A(\varphi)$. Moreover, there exists again an attraction mechanism which drives the scalar field towards a minimum of $W(\varphi)$ during the cosmological expansion. Therefore, a small value of the slope $|W_0'|$ is indeed expected at present, consistently with what is needed for solar-system tests. In conclusion, we are faced again with a serious problem: We just found a theory which seems to be consistent with all experimental data, although it is very different from GR in its field content. Our aim is therefore to find a way to distinguish it from GR, or to rule it out for internal consistency reasons.

Figure 6 displays this reconstructed coupling function $W(\varphi)$, in which the present value of the scalar field is close to the minimum at $\varphi = 0$. Its shape is nicely smooth, but its second derivative at the origin is huge if one divides it by the tiny factor $(16\pi G\hbar/c^5)H_0^2$. One gets $W_0'' \simeq 7 \times 10^{119}$, which is in fact not surprising, since the coupling function $W(\varphi)$ behaves in action (21) as the inverse of a cosmological constant. Indeed, $W(\varphi)$ multiplies the square of the curvature tensor, whereas the usual Einstein-Hilbert term involves the first

power of the scalar curvature, and a cosmological constant does not multiply any curvature term at all. It was thus expected that $W(\varphi)$ involve a dimensionless number of the order of the inverse of $(\hbar G/c^3)\Lambda \simeq 3 \times 10^{-122}$. Therefore, this model is ugly, but it is not yet ruled out. One should not confuse fine tuning and large (or small) dimensionless numbers in a model. We are here in the second situation, but there is *a priori* no fine tuning since the scalar field is attracted towards the minimum of $W(\varphi)$ during the cosmological expansion. There remains to study how efficiently it is attracted, but this is actually not necessary for our purpose.

Indeed, W_0'' takes such a gigantic value that an approximation that we made to analyze solar-system tests is no longer valid. Indeed, we have $|W_0''\varphi| \gg |W_0'|$, so that the second term on the right-hand side of Eq. (22) cannot be neglected. To simplify the discussion, we will anyway assume that $W(\varphi)$ is parabolic, which is a good approximation in a vicinity of the minimum $\varphi = 0$. We will thus neglect the higher order terms $\mathcal{O}(\varphi^2)$ in Eq. (22); taking them into account would not change our conclusions below. We did not find a close analytic solution to Eq. (22), but it is possible to write it as a series

$$\varphi = \frac{W_0'}{W_0''} \sum_{n \geq 1} \frac{1}{(3 \times 4)(7 \times 8) \cdots (4n-1)(4n)} \left(\frac{12\ell_0^2 G^2 m_\odot^2 W_0''}{r^4 c^4} \right)^n \quad (25)$$

$$\simeq \frac{W_0'}{W_0''} \left[\cos \left(\frac{Gm_\odot \ell_0}{r^2 c^2} \sqrt{3|W_0''|} \right) - 1 \right] \quad \begin{array}{l} \text{if } W_0'' < 0, \\ \text{if } W_0'' > 0. \end{array} \quad (26)$$

The second expression is a good approximation if the argument of the cosine (or hyperbolic cosine if $W_0'' > 0$) is much greater than 1. This is the case if we use the huge value of W_0'' obtained above from the cosmological reconstruction, and a typical solar-system distance for the radius r : The argument of the hyperbolic cosine is then of order 10^8 .

The above solution is such that $\varphi \propto W_0'$, therefore we do not find any nonperturbative effect similar to the “spontaneous scalarization” of neutron stars mentioned in Sec. II B above. Moreover, $\varphi \rightarrow 0$ as $r \rightarrow \infty$, and we recover GR for distances $r > 4 \times 10^{14}$ m (i.e., farther than the solar system including Oort’s comet cloud). On the other hand, there are highly nonlinear corrections proportional to $1/r^{4n}$ within the solar system. Since the ratio $(12\ell_0^2 G^2 m_\odot^2 W_0''/r^4 c^4)$ is much greater than 1, its successive powers blow up, but they are compensated by the factors $1/(3 \times 4 \times 7 \times \cdots 4n)$ which behave like the inverse of factorials. Therefore, the successive terms of series (25) start to grow exponentially, then reach a maximum for a value of the index n which may be large, and finally tend towards zero. Each of these successive terms must be assumed to be small enough for the model to pass all classical tests, but one should not forget that the largest one does not correspond to $n = 1$.

In order to study the effects of such highly nonlinear terms in the solar system, we compute their corrections to the Schwarzschild metric in the form

$$ds^2 = - \left(1 + \sum_{n \geq 1} \frac{\beta_n}{r^n} \right) c^2 dt^2 + \left(1 + \sum_{n \geq 1} \frac{\alpha_n}{r^n} \right) dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) , \quad (27)$$

and we find that the light deflection angle and the perihelion shift are respectively given by

$$\Delta\theta_* = \sum_{n \geq 1} 2^{n-1} \frac{\Gamma\left(\frac{n+1}{2}\right)^2}{\Gamma(n+1)} \frac{\alpha_n - n\beta_n}{r_0^n} + \mathcal{O}(\alpha_n, \beta_n)^2 , \quad (28)$$

$$\Delta\theta_p = \frac{6\pi Gm_\odot}{pc^2} - \sum_{n \geq 3} \frac{n(n-1)\beta_n c^2}{2Gm_\odot p^{n-1}} \pi + \mathcal{O}(\alpha_n, \beta_n)^2, \quad (29)$$

which generalize Eqs. (23)-(24) above. Note that these results are (independently) perturbative in each of the coefficients α_n and β_n , because we are assuming that the scalar-field effects are negligible with respect to the general relativistic predictions. However, the dominant scalar-field corrections may correspond to a large value of index n .

When solution (25) or its approximation (26) are used to compute the metric coefficients α_n and β_n in the above observable predictions, and if we use the huge value of W_0'' obtained from the previous cosmological reconstruction of $W(\varphi)$, we get the following experimental constraint:

$$|W_0'| < 10^{-2 \times 10^{11}}. \quad (30)$$

Now we can speak of fine tuning, and even of *hyperfine* tuning! This constraint simply means that the present value of the scalar field must be *exactly* at the minimum of the coupling function $W(\varphi)$, otherwise solar-system tests are violated. And since the universe is still evolving, the scalar field cannot remain so close to the minimum for more than a fraction of a second. Therefore, even if we assumed that $W_0' = 0$ strictly to pass solar-system tests, this would not be the case a tiny instant later. In conclusion, we managed to rule out the scalar-tensor model $A(\varphi) = 1$, $V(\varphi) = 0$ and $W(\varphi) \neq 0$. It cannot describe an accelerating expansion of the universe at present and pass solar-system (and binary-pulsar) tests at the same time.

Of course, this result does not rule out any scalar–Gauss–Bonnet coupling. A model with three (or even two) free functions $A(\varphi)$, $V(\varphi)$ and $W(\varphi)$ can obviously pass all present tests. [For instance, GR plus a cosmological constant simply corresponds to $A(\varphi) = 1$, $V(\varphi) = \Lambda/2$ and $W(\varphi) = 0$.] But the presence of a non-constant coupling $W(\varphi)$ can change the physics at small scales, notably in the very early universe (Big-Bang) and for later clustering properties.

The fact that $W(\varphi)$ induces effects at small scales can be understood by a simple dimensional argument. Since this function multiplies the square of the curvature in action (21), it induces corrections proportional to $1/r^7$ (and higher orders) to the Newtonian potential in $1/r$, and thereby generically dominates at small scales. However, we saw above that this quick reasoning can be erroneous in some perturbative but highly nonlinear situations. Indeed, if W_0'' takes very large and *negative* values, the cosine involved in Eq. (26) shows that φ is always of the order of $-W_0'/W_0''$, even for small distances r . One can then prove that the (very easily satisfied) condition $|\ell_0^2 W_0'| \ll r^2$ suffices for all scalar-field effects to be negligible in the solar system, even if $|W_0''| \sim 10^{120}$. This remark underlines that nonlinear effects can drastically change the intuitive behavior, but let us recall that our cosmological reconstruction above predicted a large and *positive* value for W_0'' . In that case, we did find that the scalar–Gauss–Bonnet coupling induces large effects at small scales, and even exponentially larger than the linear results (23)-(24).

VI. CONCLUSIONS

Scalar-tensor theories of gravity are the best motivated alternatives to general relativity. Three classes of experimental data give *qualitatively* different constraints on them. Solar-system tests strongly constrain the first derivative of the matter-scalar coupling function

$A(\varphi)$ (i.e., the linear matter-scalar coupling strength α_0). Binary-pulsar data forbid large and negative values of its second derivative β_0 (quadratic matter-scalar-scalar coupling). The knowledge of the two cosmological functions $D_L(z)$ and $\delta_m(z)$ suffices to reconstruct the full shape of both $A(\varphi)$ and the potential $V(\varphi)$ on a finite interval of φ . The knowledge of the luminosity distance $D_L(z)$ alone over a *wide* redshift interval strongly constrains the theories if one takes into account solar-system (and binary-pulsar) data, the positivity of the graviton and scalar energies, and the stability and naturalness of the models. Future data, provided by experiments like the SNAP satellite, will notably allow us to discriminate between GR plus a cosmological constant and a potential-free scalar-tensor theory. The possible coupling $W(\varphi)$ of the scalar field to the Gauss-Bonnet topological invariant can be constrained only if one takes into account cosmological and solar-system data together. The predictions of the model at small distances can depend on highly nonlinear corrections. Of course, a model including all three functions $A(\varphi)$, $V(\varphi)$ and $W(\varphi)$ is experimentally allowed, since GR plus a cosmological constant is a particular case. The presence of a scalar-Gauss-Bonnet coupling $W(\varphi)$ will generically change the behavior of the theory at small scales (clustering, Big Bang). More general tensor-multi-scalar theories, including couplings of the scalars to matter and the Gauss-Bonnet invariant, and including non-quadratic “k-essence” kinetic terms, can lead to a much richer phenomenology, but at the price of a less natural Lagrangian.

Acknowledgments

I wish to thank the organizers of the workshop for their kind invitation, and my colleagues T. Damour, D. Polarski, E. Symboloni, A.A. Starobinsky, and J.P. Uzan for collaborations or discussions on this subject.

-
- [1] G. Hinshaw, D.N. Spergel, L. Verde, *et al.*, *Astrophys. J. Suppl.* **148**, 135 (2003).
 - [2] A.G. Riess, A.V. Filippenko, P. Challis *et al.*, *Astron. J.* **116**, 1009 (1998); S. Perlmutter, G. Aldering, G. Goldhaber *et al.*, *Astroph. J.* **517**, 565 (1999).
 - [3] T. Damour and G. Esposito-Farèse, *Class. Quantum Grav.* **9**, 2093 (1992).
 - [4] P.G. Bergmann, *Int. J. Theor. Phys.* **1**, 25 (1968); K. Nordtvedt, *Astrophys. J.* **161**, 1059 (1970); R. Wagoner, *Phys. Rev. D* **1**, 3209 (1970).
 - [5] T. Damour and G. Esposito-Farèse, *Phys. Rev. Lett.* **70**, 2220 (1993); *Phys. Rev. D* **54**, 1474 (1996); **58**, 042001 (1998).
 - [6] G. Esposito-Farèse, gr-qc/0402007, to be published in the proceedings of the tenth Marcel Grossmann Meeting, July 2003; T. Damour and G. Esposito-Farèse, in preparation.
 - [7] B. Boisseau, G. Esposito-Farèse, D. Polarski, and A.A. Starobinsky, *Phys. Rev. Lett.* **85**, 2236 (2000).
 - [8] G. Esposito-Farèse and D. Polarski, *Phys. Rev. D* **63**, 063504 (2001).
 - [9] E. Symboloni, laurea thesis (2002), unpublished; G. Esposito-Farèse, gr-qc/0306018, in Proceedings of the XXXVIIIth *Rencontres de Moriond*, March 2003, Thê' Giô' Publishers (Vietnam), p. 427.
 - [10] E.G. Adelberger, B.R. Heckel, and A.E. Nelson, *Ann. Rev. Nucl. Part. Sci.* **53**, 77 (2003); C.D. Hoyle, D.J. Kapner, B.R. Heckel *et al.*, *Phys. Rev. D* **70**, 042004 (2004).

- [11] J. Khoury and A. Weltman, astro-ph/0309300; Phys. Rev. D **69**, 044026 (2004); S.S. Gubser and J. Khoury, hep-ph/0405231; P. Brax, C. van de Bruck, A.C. Davis *et al.*, astro-ph/0408415.
- [12] A.S. Eddington, *The mathematical Theory of Relativity*, Cambridge University Press (1922); K. Nordtvedt, Phys. Rev. **169**, 1017 (1968); C.M. Will and K. Nordtvedt, Astrophys. J. **177**, 757 (1972); C.M. Will, *Theory and Experiment in Gravitational Physics*, Cambridge Univ. Press (1993).
- [13] C.M. Will, *Living Rev. Rel.* **4**, 4 (2001), gr-qc/0103036.
- [14] I.I. Shapiro, in *General Relativity and Gravitation 12*, edited by N. Ashby, D.F. Bartlett, and W. Wyss, Cambridge University Press (1990), p. 313.
- [15] J.G. Williams, X.X. Newhall, and J.O. Dickey, Phys. Rev. D **53**, 6730 (1996).
- [16] T.M. Eubanks, J.O. Martin, B.A. Archinal *et al.*, Bull. Am. Phys. Soc., Abstract No. K 11.05 (1997), unpublished; draft at ftp://casa.usno.navy.mil/navnet/postscript/prd_15.ps (1999); S.S. Shapiro, J.L. Davis, D. E. Lebach, and J.S. Gregory, Phys. Rev. Lett. **92**, 121101 (2004).
- [17] B. Bertotti, L. Iess, and P. Tortora, *Nature* **425**, 374 (2003).
- [18] J.M. Weisberg and J.H. Taylor, *ASP Conference Proceedings* **302**, 93 (2003), astro-ph/0211217.
- [19] M. Bailes *et al.*, *Astrophys. J.* **595**, L49 (2003), astro-ph/0307468.
- [20] P.D. Scharre and C.M. Will. Phys. Rev. D **65**, 042002 (2002); C.M. Will and N. Yunes, Class. Quantum Grav. **21**, 4367 (2004).
- [21] T. Damour and K. Nordtvedt, Phys. Rev. D **48**, 3436 (1993).
- [22] P. Teyssandier and P. Tourrenc, J. Math. Phys. **24**, 2793 (1983).
- [23] S. Gottlöber, H.J. Schmidt, and A.A. Starobinsky, Class. Quantum Grav. **7**, 893 (1990); D. Wands, Class. Quantum Grav. **11**, 269 (1994).
- [24] K.S. Stelle, Phys. Rev. D **16**, 953 (1977); A. Hindawi, B.A. Ovrut, and D. Waldram, Phys. Rev. D **53**, 5597 (1996); E.T. Tomboulis, Phys. Lett. B **389**, 225 (1996).
- [25] T. Damour and A. Polyakov, Nucl. Phys. **B423**, 532 (1994); Gen. Rel. Grav. **26**, 1171 (1994); T. Damour, F. Piazza, and G. Veneziano, Phys. Rev. Lett. **89**, 081601 (2002); Phys. Rev. D **66**, 046007 (2002).
- [26] C. Armendariz-Picon, T. Damour, and V.F. Mukhanov, Phys. Lett. B **458**, 209 (1999); C. Armendariz-Picon, V.F. Mukhanov, and P.J. Steinhardt, Phys. Rev. D **63**, 103510 (2001); M. Malquarti, E.J. Copeland, A.R. Liddle, and M. Trodden, Phys. Rev. D **67**, 123503 (2003).